

## Safety Precautions

PPE – wear overalls, gloves, goggles

Avoid contact with eyes or skin.

Do not inhale spray mist.

If contact occurs remove contaminated clothing, rinse eyes in cold water and wash skin.

Wash hands after use

Store product in a dry cool well ventilated area.

Rinse containers before disposal

Keep out of the reach of children

Use only as an insecticide/acaricide



**DANGER**  
KEEP OUT OF REACH OF CHILDREN

# MITAMEC 1.8 EC

CONTAINS 18g/Lt ABAMECTIN

**Mitamec is a Group 6A insecticide with contact and stomach action.**

Mitamec is translaminar and partially systemic.

Mainly for the control of mites and insect pests in vegetables, soft fruit and tree crops



**PelGar International Ltd**  
Unit 5, Newman Lane, Alton, Hants.  
GU34 2QR. United Kingdom  
Tel: + 44 1420 80744  
www.pelgar.co.uk

**Distributed by:**  
PESTEX MALDIVES PVT LTD  
Pestex Building  
Neelofaru Magu  
MALE 20131  
REP OF MALDIVES

Tel: +96 03338009 / 3336878

Batch no:

Manuf. Date:

Expiry. Date:



International



Net Contents

**1 litre**

## Mixing instructions

Partly fill the spray tank and add the measured amount of **Mitamec 1.8 EC** followed by the remaining water. **Mitamec 1.8 EC** contains a built in wetter so oil additions are not required.

Use rates for control of mites, moths and leaf miners: For heavy infestations spray at 7-10 day intervals to a maximum of 2 applications per season.

CROP	MITAMEC USE RATE	WATER SPRAY VOLUME	PRE-HARVEST INTERVAL
Fruit Trees	750ml/ha	1000 l/ha	7 days citrus 14 days apple/pear
Tomato and vegetable crops	400-600ml/ha	750 l/ha	3 days
Strawberries and soft fruit	600ml/ha	Spray to the point of run off	3 days