

DESCRIPTION

Stmed-Permethrin 50EC emulsifiable concentrate has been specially developed as a repellent with long lasting activity against mosquitoes.

Stmed-Permethrin 50EC has following properties:

- A wide range of insecticidal activity, controlling both flying (resting) and crawling insects.
- A strong repellent effect against mosquitoes.
- Stmed-Permethrin 50EC has a very long lasting residual effect.
- Suitable for use as repellent on clothing and mosquitoes nets.
- Stmed-Permethrin 50EC does not accumulate in the environment.

SAFETY PRECAUTION

- Do not drink: avoid skin contact.
- Formulation operator should wear rubber gloves, rubber boots, overalls and eye protection. Stmed-Permethrin 50EC is a skin and eye irritant.
- Do not smoke, drink or eat while using this product.
- As with all chemicals, avoid contact with spray. Wear eye protection when spraying overhead or in confined places.
- Wash hand an exposed skin before meals and after work.
- Do not re-use empty containers, destroy by cutting or crushing and bury them away from habitation.
- Store in dry, cool and covered space in original container. Keep the material out of reach of children and well away from foodstuff, animal feed and food containers. Toxic to Fish and Bees.

FIRST AID

- In case of eye contact, wash thoroughly under streaming water for 10 – 15 minutes.
- In case of skin contact, remove contaminated clothing and wash affected areas with plain soap and water.
- If swallowed, give 1 – 2 glasses of water. Do not induce vomiting; take the person and this container to a doctor immediately.
- If inhaled, take the person to fresh air area.



DANGEROUS
KEEP OUT OF REACH OF CHILDREN



STMED PERMETHRIN 50EC

CONTENTS: PERMETHRIN @ 50% W/W

High performance concentrate
for space spray against
mosquito vectors.

MALARIA - DENGUE - DHF
Also controls other insect pests.

3 YEARS SHELF LIFE



Manufacturer:

 **Agropharm Ltd**
A PELGAR COMPANY

Buckingham Place, Church Road, Penn
Buckinghamshire HP10 8LN
United Kingdom
www.agropharm.co.uk
www.pelgar.co.uk

Exclusive Distributor:

STMED.,Corp

20/23 Alley 165, Thai Ha,
Dong Da, Ha noi, Viet Nam
Tel: + 84 (4) 22133555
Fax: + 84 (4) 35149865
www.stmed.com.vn

Registration No:
VNDP-HC-513-01-11

Batch no:

Manuf. Date:

Expiry. Date:



Net
Contents

1 liter

USAGE TABLE

Stmed-Permethrin 50EC can be used for residual spray, ULV spray or bed net impregnation. Stmed-Permethrin can be diluted in water or oil.

SPACE SPRAY

Application areas	Treatment	Dilution rate Stmed-Permethrin 50EC/ water, oil	Application rate
Indoors	Residual spray	25ml/ 10L	300m ² wall surface
	ULV	100ml/L	10-20ml/household
	Thermal fogging	100ml/10L	2L/ha
Outdoors	Residual spray	25ml/ 10L	300m ² wall surface
	ULV	100ml/L	0.5L/ha
	Thermal fogging	100ml/10L	2L/ha

BED NET IMPREGNATION

Type of net/size	Dilution rate Stmed-Permethrin 50EC/ water	Application rate
------------------	--	------------------

Tulle (Polyester)

- Single net (9 m ²)	3.6ml/377ml	01 single net
- Double net (16.5m ²)	6.6ml/695ml	01 double net

Cotton

- Single net (9 m ²)	3.6ml/574ml	01 single net
- Double net (16.5m ²)	6.6ml/995ml	01 double net

Please read the directions to use carefully before using

