

#### Directions for use

VULCAN AP FLY BAIT can be applied over specified fly feeding surfaces at a frequency varying from three to four weeks depending on the infestation level. VULCAN AP FLY BAIT can be used to either prevent emergence of flies or to reduce fly populations. Scatter bait from container or other convenient device. Use VULCAN AP FLY BAIT inside and around areas where flies are a nuisance. Select places where flies rest or travel (around window frames, watering places, manure storage etc.) and avoid airflows. If scattering on the floor is not possible, put the bait on a cardboard tray or plate. Scatter VULCAN AP FLY BAIT (do not put in piles) at the rate of 25g per 10 square meters of fly feeding area. Put the bait stations out of the reach of animals. This product should not be used continuously against houseflies in intensive or controlled environment units as this could cause control failure due to insecticide resistance. If sequential treatments are required, use a product with a different active substance and, if necessary, a different control method (such as a space spray). In any one location, any neonicotinoid containing product (and including any control method) must not be used more than 6 times in any 12 month period.

#### PelGar International Ltd.

Unit 13 Newman Lane,  
Alton. GU34 2QR.  
Tel: 01420 80744  
Email: info@pelgar.co.uk



Authorisation holder: Kwizda, France.



International

# VULCAN AP FLY BAIT

Acetamiprid 0.495% w/w (CAS No. 160430-64-8)  
Contains the bittering agent (denatonium benzoate)

**VULCAN AP FLY BAIT is a granular bait insecticide for the control of flies in a wide variety of breeding environments (e.g. cattle, pigs, poultry, sheep) and very easy to use. VULCAN AP FLY BAIT has an excellent attractiveness towards flies with the dual effect of a food attractant and a sexual pheromone.**



# 500g

#### Precautions

Keep out of reach of children. Call a POISON CENTER or doctor/physician if you feel unwell. Dispose of contents/container in accordance with local requirements. When using, do not eat, drink or smoke. Do not re-use empty contaminated packaging for other purposes. The COSHH (Control of Substances Hazardous to Health) Regulations 2002 (as amended) apply to the use of this product at work. This material and its container must be disposed of in a safe way. Avoid excessive contamination of coveralls. DO NOT USE where food, feed or water could become contaminated. PREVENT ACCESS TO BAITS by children and animals. DO NOT CONTAMINATE WATER WITH PRODUCT OR CONTAMINATED MANURE/SLURRY. AVOID CONTAMINATION VIA DRAINS FROM FARMYARDS AND ANIMAL HOUSING/BREEDING UNITS. ALL MANURE/SLURRY FROM THE PREMISES SHOULD BE COLLECTED AND DISPOSED OF VIA LAND FILL ACCORDING TO LOCAL LEGISLATION. HAZARDOUS TO BEES. COVER ALL WATER STORAGE TANKS before application. DO NOT APPLY DIRECTLY TO ANIMALS. EXCLUDE ANIMALS DURING APPLICATION. Protect milk machinery and milk containers from contamination. COLLECT EGGS AND REMOVE FEED, EXPOSED WATER AND MILK BEFORE APPLICATION. WASH HANDS AND EXPOSED SKIN before meals and after use. KEEP IN A SAFE PLACE.

#### STATUTORY CONDITIONS OF APPROVAL RELATING TO USE.

INSECTICIDE FOR PROFESSIONAL USE AGAINST FLIES. FOR USE IN BUILDINGS WHERE ANIMALS ARE HOUSED FOR BREEDING. SCATTER AT THE RATE OF 25g PER 10 SQUARE METRES OF FLY FEEDING AREA.

Users must comply in full with all their legal responsibilities under the Control of Pesticides Regulations 1986 (as amended), including complying with all Conditions of Approval.

**READ ALL PRECAUTIONS BEFORE USE.**

**HSE No. 9827**

To avoid risks to human health and the environment, comply with the instructions for use.

Batch number: See pack.

DOE: See pack.