

WARNING - KEEP OUT OF REACH OF CHILDREN

Statement of practical treatment of poisoning.

IF SWALLOWED do not induce vomiting. Administer entire contents of bottle of Carbomix.

Perform gastric lavage under medical supervision and re-administer Carbomix.

IF INHALED move patient to fresh air and summon medical assistance if signs of respiratory distress are evident.

IF ON SKIN remove contaminated clothing and wash with soap and water.

IF IN EYES flush with water for at least 15 minutes and obtain medical assistance if symptoms persist.

WARNING

HARMFUL IF SWALLOWED

FOR USE ONLY AS AN INSECTICIDE

FLAMMABLE

Agromethrin is toxic to aquatic animals and is harmful to bees.

STORAGE AND DISPOSAL

Store in original container in a cool place and out of reach of children.

Thoroughly wash out container and dispose of contents according to local laws. Do not reuse container. Bury in non crop area away from water.

Observe all regulations regarding handling of pesticides.



ANGER
KEEP OUT OF REACH OF CHILDREN

TROPIVITE® AGROMETHRIN 40 EC

CONTAINS 40g/Lt ALPHA CYPERMETHRIN

Alpha-cyano-3-phenoxybenzyl cis-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropane-carboxylate
Solvent system (xylene & aliphatics) emulsifiers and wetting agents to 100%



PelGar International Ltd
Unit 5, Newman Lane, Alton, Hants.
GU34 2QR. United Kingdom
Tel: + 44 1420 80744
www.pelgar.co.uk

Distributed by:
PESTEX MALDIVES PVT LTD
Pestex Building
Neelofaru Magu
MALE 20131
REP OF MALDIVES

Tel: +96 03338009 / 3336878

Batch no:

Manuf. Date:

Expiry. Date:



International



Net Contents

1 litre

Recommendations

Apply using tractor sprayer or knapsack in dilution of 1-500 or 1-1000 in clean water. For aerial application dilute 1-30 or 1-50. Apply at rate of 1 litre of concentrate per hectare.

Properties and Use rates:

Contact emulsifiable insecticide used to control a wide range of sucking and biting insects.

Agromethrin is an insecticide of low toxicity and operates at a low concentration to control insects that attack agricultural crops.

CROPS	INSECTS	USE RATE AGROMETHRIN/100L WATER
Field crops: Alfalfa, clover, Cotton & Rhodes Corn	Weevils, aphids, leaf miner, biting insects, leaf worms. Cotton insects and cotton acanthoid worm.	50-100ml 150-225ml
Vegetable crops: Tomato, cucumber, onion, potato, cabbage, carrot, eggplant, watermelon	White flies, aphids, thrips, fruit worm, biting insects and leaf miner.	50-100ml
Fruit crops: Grape, apple, citrus, olive, mango, peach, palm trees	Aphids, thrips, fruit worms, scales, weevils, mealybugs and leaf miner	50-100ml

SAFETY PERIOD : 5 Days

Ensure complete coverage of target crop at first sign of infestation and repeat as necessary